PARKLAND MEMORIAL HOSPITAL

PATHOLOGY LABORATORY

Mudertolder BLB Autopsy Number: N53-352 Zule

Name: Tippit, J. D.

Admitted: 11-22-63 DOA

Autopsy date: 11-22-63, 3:15 F.M.

Autopsy by: Earl F. Rose, M. D.

Hone Restrictions:

Unit No.

Sex: Male Age: 39 Race: White

Expired:

Service:

Misight Coroner: Judge Joe B. Brown, Jr.

external examination: 🕾

External examination reveals a well developed white male measuring 5 foot, 11 inches in length and weighing an estimated 175 to 180 pounds. The hair is black, slightly very, very slight frontal balding. The irides are blue, the pupils are equal at 5 mm. Very slight posterior mottled lividity and body heat is present. Rigor is not present. The neck is not remrisable. Hair distribution is normal, the Oral hygiene is good. Identification bands on the left wrist, right wrist and left penis is circurcised. ankle.

There are four entrance types of wounds. No. 1 is 4 3/4 inches from the top of the head end 3 3/4 inches to the right of the midline. This measures $3/8 \times 1/4$ inch and is corremaed by a contusion ring. He powder tettooing is noted at the margins.

Wound No. 2 is 17 inches from the top of the head on the right chest. It is 4 inches to the right of the midline, above and slightly medial to the right nipple. It measures 3/8 x 1/4 of an inch, surramded by bruising. There is also a contusion ring.

tound No. 3 is 21 inches from the top of the head, along the exterior lateral side of the right chest and is 6 inches to the right of the midline. This measures 5/16 x 1/4 of an inch and is surrounded by a contusion ring.

Vound No. 4 on the left chest is 20% inches from the top of the head, 1% inches to the left of the midline. The wound measures 3/4 x 3/8 of an inch, is transverse and surrounding this is a 1/4 x 3/4 inch abrasion.

There is taxaing of the arms. On the left arm there is a tattoo being "Tippit". On the dorson of the left hand there is a crusted chroded area acasuring à x à inch and a fresh abraded area on the dorsum of the right hand which measures i x i inch. The nails are quite well cared for elthough slightly dirty. Scar above the left bace, runs in an oblique fachion, crosses to the mainl espect of the knee, terminates on the leg measuring 7 inches. Poorly defined i inch impoulation type of scar on the left deltoid region.

16) inches from the top of the head, 3/4 of an inch to the right of the midline of the back in the subcutaneous tiesus a misello is recovered froz this region. This is associated with the Wound No. 3.

INCISIONS: The standard "I" thoraconbiominal and intermestaid incisions are utilized. Examination of the wound of the right temple is made. It is found to enter in the right middle cranial force, pursues a course which is elightly upward, backward and