

cut 2 cm. transverse incision into the
subcutaneous tissue.

There is an old, well healed 8 cm.
Bunny abdominal incision. On the
umbilicus the midline is an old, well
healed 15 cm. scar. Situated on the
upper anterior-lateral aspect of the st.
thigh is an old, well healed 8 cm. scar.

Skull Wounds

There is a large irregular defect of the
skull and scalp on the right involving
chiefly the parietal bone but extending
slightly into the temporal and occipital
regions. In this region there is an
total absence of scalp and bone producing
defect which measures approximately
4 cm. in greatest diameter.

From the irregular margins of the
bone scalp defect tears extend in stellate
fashion into more or less intact scalp
as follows:

a) From the right inferior temporo-

Exhibit 38

parietal margin anterior to the st. ear to
a point slightly above the tragus.

b) From the anterior parietal margin
anteriorly on the forehead to approximately
4 cm. above the st. orbital ridge.

c) From the left margin of the main
defect across the midline anterior-laterally
for a distance of approximately 8 cm.

d) From the same starting point as
c) 10 cm. posterior-laterally.

Situated in the posterior scalp approximately
2.5 cm. laterally to the right and slightly
above the external occipital protuberance
is a ~~punctured~~ ^{lacerated} wound ~~tangential to the~~
~~surface of the scalp~~ measuring 15 x 6 mm.

In the underlying bone is a corresponding
~~puncture~~ ^{puncture} wound through ~~the bone~~
the skull which exhibits ~~irregular~~ ^{irregular}
the margins of the bone when ~~punctured~~ ^{viewed}
the inner ~~side~~ ^{aspect} of the skull.

A clearly visible in the above described
large skull defect and extending from it
is ~~lacerated~~ ^{lacerated} brain tissue which on
close inspection proves to represent ~~the~~

Humes twice omitted the word "puncture" on this page, once actually replacing it with
a word of entirely opposite meaning, "lacerated."